

# The Volterra lattice as a gradient flow

A. V. Penskoï\*

## Abstract

The Volterra lattice is considered. New gradient interpretation for this dynamical system is proposed. This interpretation seems to be more natural than existing ones.

**AMS MSC 58F19, 58F07**

In the present paper we give a gradient interpretation for the Volterra lattice treated in details in [1]. The phase space consists of variables  $u_n > 0$ ,  $1 \leq n \leq N$ . The equations of motion are

$$\frac{du_n}{dt} = u_n(u_{n+1} - u_{n-1}),$$

where  $u_0 = u_{N+1} = 0$ .

The interest to gradient interpretations goes back to the paper by J. Moser [2], where such an interpretation was given implicitly for the Toda lattice. Twenty years later A. Bloch, R. Brockett and T. Ratiu [3] suggested another gradient interpretation for the Toda lattice, which was different from Moser's one. Our interpretation for the Volterra lattice as a gradient flow is based on the same idea of the double bracket representation (see [3]).

It should be noted that as well as for the Toda lattice our interpretation is not unique: the well-known isomorphism between these two problems (see e.g. [4]) and the results of [3] lead to one more gradient interpretation for the Volterra lattice. Our representation seems to be more natural. As a remark we note that the gradient interpretation for the Toda lattice was used in the paper by C. Tomei [5] and D. Fried [6] for study of non-trivial geometry of isospectral sets.

---

\*Faculty of Mechanics and Mathematics, Department of Higher Geometry and Topology, Moscow State University, Vorob'ievsky Gory, Moscow, 119899, Russia. e-mail: penskoi@mech.math.msu.su

Let  $c_i = \sqrt{u_i}$ . Let us consider a manifold  $M$  of all matrices of the form

$$L = \begin{pmatrix} 0 & c_1 & 0 & \dots & 0 \\ c_1 & 0 & c_2 & & \vdots \\ 0 & c_2 & 0 & \ddots & \vdots \\ \vdots & & \ddots & 0 & c_N \\ 0 & \dots & \dots & c_N & 0 \end{pmatrix}$$

with a given spectrum. The manifold  $M$  is called the isospectral set. Let  $V$  be the euclidean space of  $(N+1) \times (N+1)$ -matrices with respect to the scalar product  $\langle X, Y \rangle = \text{tr}XY^T$ . The manifold  $M$  is naturally embedded in  $V$ . Let  $L_0$  be any point of  $M$ . Consider the adjoint  $GL_{N+1}$ -action in  $V$ ; then  $GL_{N+1}L_0$  is a submanifold of  $V$  and  $M$  is a submanifold of  $GL_{N+1}L_0$ . Let  $V_{L_0} = \{T \in V | [L_0, T] = 0\}$ , let  $V_{L_0}^\perp$  be its orthogonal complement with respect to  $\langle, \rangle$ ; then  $V_{L_0}^\perp \cong T_{L_0}GL_{N+1}L_0$ . By  $T^\perp$  denote the orthogonal projection of  $T \in V$  on  $V_{L_0}^\perp$ . Define the scalar product of two vectors  $[L_0, A], [L_0, B] \in T_{L_0}GL_{N+1}L_0$  as  $([L_0, A], [L_0, B]) = \langle A^\perp, B^\perp \rangle$ . In this way we obtain a Riemannian metric on  $GL_{N+1}L_0$ . Its restriction gives a Riemannian metric on  $M$ .

Let  $V \ni K = \frac{1}{4}\text{diag}(1, 2, 3, 4, \dots)$  and  $f(L) = \langle K, L^2 \rangle = \text{tr}KL^2$ .

**THEOREM.** The gradient flow of  $f|_M$  relative to the Riemannian metric on  $M$  coincides with the restriction of Volterra flow to a joint level of its integrals.

**PROOF.** It is known (see e.g. [4]) that the Volterra lattice is equivalent to the Lax equation  $\dot{L} = [L, A]$ , where  $A$  is the matrix

$$\begin{pmatrix} 0 & 0 & \frac{1}{2}c_1c_2 & 0 & \dots & 0 \\ 0 & 0 & 0 & \frac{1}{2}c_2c_3 & & \vdots \\ -\frac{1}{2}c_1c_2 & 0 & 0 & & \ddots & \\ 0 & -\frac{1}{2}c_2c_3 & & & & 0 \\ \vdots & & \ddots & & & \frac{1}{2}c_{N-1}c_N \\ 0 & \dots & 0 & -\frac{1}{2}c_{N-1}c_N & 0 & 0 \end{pmatrix}.$$

The integrals of motion are  $\text{tr}L^k$ ; therefore  $M$  is diffeomorphic to the joint level of integrals. It is sufficient to prove that the restriction on  $M$  of the gradient flow of  $f|_{GL_{N+1}L}$  relative to the Riemannian metric  $(,)$  coincides with the Volterra flow.

Let  $[L, T] \in T_L GL_{N+1} L$  be an arbitrary vector. Let us consider  $X$  such that  $[L, X] = \text{grad}(f|_{GL_{N+1} L})$ ; then we have the equation  $df([L, T]) = ([L, X], [L, T])$ . We see that  $df([L, T]) = \langle K, L[L, T] + [L, T]L \rangle = \langle K, [L^2, T] \rangle = \langle [L^2, K], T \rangle$ . Further,  $[L^2, K]^\perp = [L^2, K]$ , therefore  $df([L, T]) = \langle [L^2, K]^\perp, T \rangle$ . At the same time  $([L, X], [L, T]) = \langle X^\perp, T^\perp \rangle$ ; then  $X^\perp = [L^2, K]$  and  $\text{grad}(f|_{GL_{N+1} L}) = [L, [L^2, K]]$ .

On the other hand it is easy to prove that  $A = [L^2, K]$ , thus the Volterra lattice is equivalent to the equation  $\dot{L} = [L, [L^2, K]]$ . This completes the proof.  $\square$

## References

- [1] FADEEV L. D., TAKHTAJAN L. A. Hamiltonian methods in the theory of solitons. Springer, Berlin, 1986.
- [2] MOSER J. Finitely many mass points on the line under the influence of an exponential potential — an integrable system. // Lecture Notes in Phys., 1975, V.38, P. 467-497.
- [3] BLOCH A. M., BROCKETT R. W., RATIU T.S. Completely Integrable Gradient Flows // Comm. Math. Phys. 1992. V. 147. N 1. P. 57-74.
- [4] VERESHCHAGIN V. L. Spectral theory of monophase solutions of the Volterra lattice // Mat. Zametki 1990. V. 48. N 2. P. 145-148.
- [5] TOMEI C. The topology of isospectral manifolds of tridiagonal matrices. // Duke Math. J. 1984. V. 51. N 4. P. 981-996.
- [6] FRIED D. The cohomology of an isospectral flow. // Proc. Amer. Math. Soc. 1986. V. 98. N 2. P. 363-368.